Complete Summary

TITLE

Immunizations: percentage of adolescents who are up-to-date with recommended immunizations (Hep B, MMR, tetanus, and verification of varicella immunity).

SOURCE(S)

Institute for Clinical Systems Improvement (ICSI). Immunizations. Bloomington (MN): Institute for Clinical Systems Improvement (ICSI); 2004 Jun. 56 p. [61 references]

Brief Abstract

DESCRIPTION

This measure assesses the percentage of adolescents who are up-to-date with recommended immunizations (Hep B, MMR, tetanus, and verification of varicella immunity).

RATIONALE

The priority aim addressed by this measure is to increase the rate of people upto-date with recommended immunizations.

PRIMARY CLINICAL COMPONENT

Immunization; Hepatitis B (Hep B); measles, mumps, rubella (MMR); tetanus and diphtheria toxoids (Td); varicella (VZV)

DENOMINATOR DESCRIPTION

All adolescents age 12 to 17 with continuous enrollment who presented for a primary care visit in the specified quarter

NUMERATOR DESCRIPTION

Number of adolescents in denominator who are up-to-date with recommended immunizations (see the related "Numerator Inclusions/Exclusions" field the Complete Summary)

Evidence Supporting the Measure

PRIMARY MEASURE DOMAIN

Process

SECONDARY MEASURE DOMAIN

Not applicable

EVIDENCE SUPPORTING THE MEASURE

A clinical practice guideline or other peer-reviewed synthesis of the clinical evidence

NATIONAL GUIDELINE CLEARINGHOUSE LINK

• <u>Immunizations.</u>

Evidence Supporting Need for the Measure

NEED FOR THE MEASURE

Unspecified

State of Use of the Measure

STATE OF USE

Current routine use

CURRENT USE

Internal quality improvement

Application of Measure in its Current Use

CARE SETTING

Physician Group Practices/Clinics

PROFESSIONALS RESPONSIBLE FOR HEALTH CARE

Physicians

LOWEST LEVEL OF HEALTH CARE DELIVERY ADDRESSED

Group Clinical Practices

TARGET POPULATION AGE

Age 12 to 17 years

TARGET POPULATION GENDER

Either male or female

STRATIFICATION BY VULNERABLE POPULATIONS

Unspecified

Characteristics of the Primary Clinical Component

INCIDENCE/PREVALENCE

For immunized children with household exposure to varicella, the attack rate is 25%. For unimmunized children with household exposure, the rate is 90%. This is an effectiveness rate of 72%. However, the vaccine is more than 95% effective in preventing serious disease (greater than 300 lesions and a fever). With natural exposure (i.e., just being in the community) a 1 to 3% annual breakthrough rate occurs in vaccinated children.

EVIDENCE FOR INCIDENCE/PREVALENCE

Institute for Clinical Systems Improvement (ICSI). Immunizations. Bloomington (MN): Institute for Clinical Systems Improvement (ICSI); 2004 Jun. 56 p. [61 references]

ASSOCIATION WITH VULNERABLE POPULATIONS

Unspecified

BURDEN OF ILLNESS

Unspecified

UTILIZATION

Unspecified

COSTS

Unspecified

Institute of Medicine National Healthcare Quality Report Categories

IOM CARE NEED

Staying Healthy

IOM DOMAIN

Data Collection for the Measure

CASE FINDING

Users of care only

DESCRIPTION OF CASE FINDING

Adolescents ages 12 to 17

This measure should be collected through:

- Identifying eligible adolescents who presented in the target quarter for a primary care visit who have maintained continuous enrollment for 18 months.
- Reviewing the record to determine if all recommended doses were given within the recommended time frames.

The suggested time period for data collection is quarterly.

DENOMINATOR SAMPLING FRAME

Patients associated with provider

DENOMINATOR (INDEX) EVENT

Encounter Patient Characteristic

DENOMINATOR INCLUSIONS/EXCLUSIONS

Inclusions

All adolescents age 12 to 17 with continuous enrollment who presented for a primary care visit in the specified quarter

Exclusions Unspecified

NUMERATOR INCLUSIONS/EXCLUSIONS

Inclusions

Number of adolescents in denominator who are up-to-date with recommended immunizations*

- Hep B:
 - 3 doses (all given within 18 months of the first Hep B dose)

^{*}An adolescent is up-to-date with recommended immunizations if:

- 2 doses Recombivax® HB given 4 to 6 months apart (see "Notes" below)
- MMR: 2 doses (first dose greater than or equal to 1 yr old; second dose 4 to 6 years)
- Td: 3 or 4 doses (second dose 4 weeks after first; third dose 6 months after second dose; booster every 10 years)
- VZV: 1 dose or documented history of varicella

Notes: Adolescents (age 11 to 15 years) can be immunized with Recombivax® HB per an alternative 2-dose schedule of 10 micrograms intramuscularly (IM) given 4 to 6 months apart. Be aware, however, that the introduction of this alternate dosing schedule does increase operational complexity for immunization administration in a clinic serving multiple ages of children and will therefore increase resultant risk of error.

Exclusions Unspecified

DENOMINATOR TIME WINDOW

Time window is a single point in time

NUMERATOR TIME WINDOW

Fixed time period

DATA SOURCE

Administrative data Medical record

LEVEL OF DETERMINATION OF QUALITY

Individual Case

PRE-EXISTING INSTRUMENT USED

Unspecified

Computation of the Measure

SCORING

Rate

INTERPRETATION OF SCORE

Better quality is associated with a higher score

ALLOWANCE FOR PATIENT FACTORS

Unspecified

STANDARD OF COMPARISON

Internal time comparison

Evaluation of Measure Properties

EXTENT OF MEASURE TESTING

Unspecified

Identifying Information

ORIGINAL TITLE

Percentage of adolescents who are up-to-date with recommended immunizations (Hep B, MMR, tetanus, and verification of varicella immunity).

MEASURE COLLECTION

<u>Immunizations Measures</u>

DEVELOPER

Institute for Clinical Systems Improvement

ADAPTATION

Measure was not adapted from another source.

RELEASE DATE

2003 Jul

REVISION DATE

2004 Jun

MEASURE STATUS

This is the current release of the measure.

This measure updates a previous version: Institute for Clinical Systems Improvement (ICSI). Immunizations. Bloomington (MN): Institute for Clinical Systems Improvement (ICSI); 2003 Jul. 51 p.

SOURCE(S)

Institute for Clinical Systems Improvement (ICSI). Immunizations. Bloomington (MN): Institute for Clinical Systems Improvement (ICSI); 2004 Jun. 56 p. [61 references]

MEASURE AVAILABILITY

The individual measure, "Percentage of adolescents who are up-to-date with recommended immunizations (Hep B, MMR, tetanus, and verification of varicella immunity)," is published in "Health Care Guideline: Immunizations." This document is available from the Institute for Clinical Systems Improvement (ICSI) Web site.

For more information, contact ICSI at, 8009 34th Avenue South, Suite 1200, Bloomington, MN 55425; phone: 952-814-7060; fax: 952-858-9675; Web site: www.icsi.org; e-mail: icsi.info@icsi.org.

NQMC STATUS

This NQMC summary was completed by ECRI on April 19, 2004. This summary was updated by ECRI on October 11, 2004.

COPYRIGHT STATEMENT

This NQMC summary (abstracted Institute for Clinical Systems Improvement [ICSI] Measure) is based on the original measure, which is subject to the measure developer's copyright restrictions.

The abstracted ICSI Measures contained in this Web site may be downloaded by any individual or organization. If the abstracted ICSI Measures are downloaded by an individual, the individual may not distribute copies to third parties.

If the abstracted ICSI Measures are downloaded by an organization, copies may be distributed to the organization's employees but may not be distributed outside of the organization without the prior written consent of the Institute for Clinical Systems Improvement, Inc.

All other copyright rights in the abstracted ICSI Measures are reserved by the Institute for Clinical Systems Improvement, Inc. The Institute for Clinical Systems Improvement, Inc. assumes no liability for any adaptations or revisions or modifications made to the abstracts of the ICSI Measures.

© 2004 National Quality Measures Clearinghouse

Date Modified: 11/1/2004

FirstGov

